

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Book	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			

Entrez PubMed

Display	Summary	▼	Sort	▼	Save	Text	Add to Clipboard	C						
Show:	20	▼	Items 1-20 of 262	Page 1 of 14	Select page:	1	2	3	4	5	6	7	8	9

PubMed Services

- ☐ 1: [Scholl FA, McLoughlin P, Ehler E, de Giovanni C, Schafer BW.](#) Related Articles, Nucl
DRAL is a p53-responsive gene whose four and a half LIM domain protein pro
induces apoptosis.
J Cell Biol. 2000 Oct 30;151(3):495-506.
PMID: 11062252 [PubMed - indexed for MEDLINE]

Related Resources

- ☐ 2: [Zeng X, Keller D, Wu L, Lu H.](#) Related /
UV but not gamma irradiation accelerates p53-induced apoptosis of teratocarci
cells by repressing MDM2 transcription.
Cancer Res. 2000 Nov 1;60(21):6184-8.
PMID: 11085543 [PubMed - indexed for MEDLINE]

- ☐ 3: [Juven T, Barak Y, Zauberman A, George DL, Oren M.](#) Related Articles, Nucl
Wild type p53 can mediate sequence-specific transactivation of an internal pron
within the mdm2 gene.
Oncogene. 1993 Dec;8(12):3411-6.
PMID: 8247544 [PubMed - indexed for MEDLINE]

- ☐ 4: [Janssens V, Van Hoof C, De Baere I, Merlevede W, Goris J.](#) Related /
The phosphotyrosyl phosphatase activator gene is a novel p53 target gene.
J Biol Chem. 2000 Jul 7;275(27):20488-95.
PMID: 10787423 [PubMed - indexed for MEDLINE]

- ☐ 5: [Shiraishi K, Fukuda S, Mori T, Matsuda K, Yamaguchi T, Tanikawa C, Ogawa M, Nakamura Y, Arakawa H.](#) Related Articles,
Identification of fractalkine, a CX3C-type chemokine, as a direct target of p53.
Cancer Res. 2000 Jul 15;60(14):3722-6.
PMID: 10919640 [PubMed - indexed for MEDLINE]

- ☐ 6: [Kostic C, Shaw PH.](#) Related /
Isolation and characterization of sixteen novel p53 response genes.
Oncogene. 2000 Aug 17;19(35):3978-87.
PMID: 10962554 [PubMed - indexed for MEDLINE]


- ☐ 7: [Theis S, Atz J, Mueller-Lantzsch N, Roemer K.](#) Related /
A function in apoptosis other than transactivation inherent in the NH2-terminal
domain of p53.
Int J Cancer. 1997 May 29;71(5):858-66.
PMID: 9180157 [PubMed - indexed for MEDLINE]

- ☐ **8:** [Cervellera M, Raschella G, Santilli G, Tanno B, Ventura A, Mancini C, Seignani C, Calabretta B, Sala A.](#) Related /
Direct transactivation of the anti-apoptotic gene apolipoprotein J (clusterin) by B-MYB.
J Biol Chem. 2000 Jul 14;275(28):21055-60.
PMID: 10770937 [PubMed - indexed for MEDLINE]
- ☐ **9:** [Venot C, Maratrat M, Sierra V, Conseiller E, Debussche L.](#) Related /
Definition of a p53 transactivation function-deficient mutant and characterization of two independent p53 transactivation subdomains.
Oncogene. 1999 Apr 8;18(14):2405-10.
PMID: 10327062 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Resnick-Silverman L, St Clair S, Maurer M, Zhao K, Manfredi JJ.](#) Related /
Identification of a novel class of genomic DNA-binding sites suggests a mechanism for selectivity in target gene activation by the tumor suppressor protein p53.
Genes Dev. 1998 Jul 15;12(14):2102-7.
PMID: 9679054 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Roth J, Koch P, Contente A, Dobbstein M.](#) Related /
Tumor-derived mutations within the DNA-binding domain of p53 that phenotypically resemble the deletion of the proline-rich domain.
Oncogene. 2000 Mar 30;19(14):1834-42.
PMID: 10777217 [PubMed - indexed for MEDLINE]
- ☐ **12:** [Genini M, Schwalbe P, Scholl FA, Remppis A, Mattei MG, Schafer BW.](#) Related Articles, Nucleotide, OMIM, F
Subtractive cloning and characterization of DRAL, a novel LIM-domain protein down-regulated in rhabdomyosarcoma.
DNA Cell Biol. 1997 Apr;16(4):433-42.
PMID: 9150430 [PubMed - indexed for MEDLINE]
- ☐ **13:** [Sutcliffe T, Fu L, Abraham J, Vaziri H, Benchimol S.](#) Related /
A functional wild-type p53 gene is expressed in human acute myeloid leukemia lines.
Blood. 1998 Oct 15;92(8):2977-9. No abstract available.
PMID: 9763589 [PubMed - indexed for MEDLINE]
- ☐ **14:** [Yang X, Pater A, Tang SC.](#) Related Articles,
Cloning and characterization of the human BAG-1 gene promoter: upregulation of tumor-derived p53 mutants.
Oncogene. 1999 Aug 12;18(32):4546-53.
PMID: 10467399 [PubMed - indexed for MEDLINE]
- ☐ **15:** [Deguin-Chambon V, Vacher M, Jullien M, May E, Bourdon JC.](#) Related /
Direct transactivation of c-Ha-Ras gene by p53: evidence for its involvement in transactivation activity and p53-mediated apoptosis.
Oncogene. 2000 Nov 30;19(51):5831-41.
PMID: 11127813 [PubMed - indexed for MEDLINE]

- ☐ **16:** [Schafer H, Trauzold A, Sebens T, Deppert W, Folsch UR, Schmidt WE.](#) [Related /](#)
The proliferation-associated early response gene p22/PRG1 is a novel p53 target gene.
Oncogene. 1998 May 14;16(19):2479-87.
PMID: 9627114 [PubMed - indexed for MEDLINE]
- ☐ **17:** [Tan M, Heizmann CW, Guan K, Schafer BW, Sun Y.](#) [Related /](#)
Transcriptional activation of the human S100A2 promoter by wild-type p53.
FEBS Lett. 1999 Feb 26;445(2-3):265-8.
PMID: 10094469 [PubMed - indexed for MEDLINE]
- ☐ **18:** [Xu D, Wang Q, Gruber A, Bjorkholm M, Chen Z, Zaid A, Selivanova G, Peterson C, Wiman KG, Pisa P.](#) [Related /](#)
Downregulation of telomerase reverse transcriptase mRNA expression by wild p53 in human tumor cells.
Oncogene. 2000 Oct 26;19(45):5123-33.
PMID: 11064449 [PubMed - indexed for MEDLINE]
- ☐ **19:** [Gottlieb E, Lindner S, Oren M.](#) [Related /](#)
Relationship of sequence-specific transactivation and p53-regulated apoptosis in interleukin 3-dependent hematopoietic cells.
Cell Growth Differ. 1996 Mar;7(3):301-10.
PMID: 8838860 [PubMed - indexed for MEDLINE]
- ☐ **20:** [Selivanova G, Iotsova V, Okan I, Fritsche M, Strom M, Groner B, Grafstrom RC, Wiman KG.](#) [Related /](#)
Restoration of the growth suppression function of mutant p53 by a synthetic peptide derived from the p53 C-terminal domain.
Nat Med. 1997 Jun;3(6):632-8.
PMID: 9176489 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Sort	▼	Save	Text	Add to Clipboard	C						
Show:	20	▼	Items 1-20 of 262	Page 1 of 14	Select page:	1	2	3	4	5	6	7	8	9

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Book	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			
Display		Summary	▼	Sort	▼	Save	Text	Add to Clipboard	
Show: 20		▼	Items 1-20 of 105		Page 1 of 6		Select page: 1 2 3		

Entrez PubMed

PubMed Services

Related Resources

- ☐ 1: Genini M, Schwalbe P, Scholl FA, Remppis A, Mattei MG, Schafer BW. Related Articles, Nucleotide, OMIM, F
Subtractive cloning and characterization of DRAL, a novel LIM-domain protein down-regulated in rhabdomyosarcoma.
DNA Cell Biol. 1997 Apr;16(4):433-42.
PMID: 9150430 [PubMed - indexed for MEDLINE]
- ☐ 2: Russell MW, Baker P, Izumo S. Related Articles, Nucleotide, F
Cloning, chromosomal mapping, and expression of the human eHAND gene.
Mamm Genome. 1997;8(11):863-5. No abstract available.
PMID: 9337404 [PubMed - indexed for MEDLINE]
- ☐ 3: Racevskis J, Dill A, Sparano JA, Ruan H. Related Articles, Nucleotide, F
Molecular cloning of LMO41, a new human LIM domain gene.
Biochim Biophys Acta. 1999 Apr 14;1445(1):148-53.
PMID: 10209267 [PubMed - indexed for MEDLINE]
- ☐ 4: Kibardin AV, Mirkina II, Korneeva EA, Gnuchev NV, Georgiev GP, Kiselev SL. Related /
Molecular cloning of a new mouse gene tagL containing a lysozyme-like domain.
Dokl Biochem. 2000 May-Jun;372(1-6):103-5. No abstract available.
PMID: 10935177 [PubMed - indexed for MEDLINE]
- ☐ 5: Chan KK, Tsui SK, Lee SM, Luk SC, Liew CC, Fung KP, Waye MM, Lee CY. Related Articles, Nucleotide, OMIM, F
Molecular cloning and characterization of FHL2, a novel LIM domain protein preferentially expressed in human heart.
Gene. 1998 Apr 14;210(2):345-50.
PMID: 9573400 [PubMed - indexed for MEDLINE]
- ☐ 6: Nambiar A, Kandala JC, Svoboda J, Guntaka RV. Related /
Cloning of a novel Y-box homology protein (chkYB-1HP) cDNA lacking the cold-shock domain.
Biochim Biophys Acta. 1998 Jan 7;1395(1):1-6.
PMID: 9434143 [PubMed - indexed for MEDLINE]
- ☐ 7: Vidal-Taboada JM, Lu A, Pique M, Pons G, Gil J, Oliva R. Related Articles, Nucleotide, OMIM, F
Down syndrome critical region gene 2: expression during mouse development and human cell lines indicates a function related to cell proliferation.

Biochem Biophys Res Commun. 2000 May 27;272(1):156-63.
PMID: 10872820 [PubMed - indexed for MEDLINE]

- ☐ **8:** Mason P, Bayol S, Brownson C, Loughna P. Related /
Selective expression of a novel striated muscle protein.
Biochem Soc Trans. 1998 May;26(2):S138. No abstract available.
PMID: 9649813 [PubMed - indexed for MEDLINE]
- ☐ **9:** Nagasaki K, Manabe T, Hanzawa H, Maass N, Tsukada T, Yamaguchi K. Related Articles, Nucleotide, F
Identification of a novel gene, LDOC1, down-regulated in cancer cell lines.
Cancer Lett. 1999 Jun 1;140(1-2):227-34.
PMID: 10403563 [PubMed - indexed for MEDLINE]
- ☐ **10:** Genini M, Schwalbe P, Scholl FA, Schafer BW. Related Articles, Nucl
Isolation of genes differentially expressed in human primary myoblasts and embryonal rhabdomyosarcoma.
Int J Cancer. 1996 May 16;66(4):571-7.
PMID: 8635876 [PubMed - indexed for MEDLINE]
- ☐ **11:** Scholl FA, McLoughlin P, Ehler E, de Giovanni C, Schafer BW. Related Articles, Nucl
DRAL is a p53-responsive gene whose four and a half LIM domain protein product induces apoptosis.
J Cell Biol. 2000 Oct 30;151(3):495-506.
PMID: 11062252 [PubMed - indexed for MEDLINE]
- ☐ **12:** Ozaki K, Nagata M, Suzuki M, Fujiwara T, Miyoshi Y, Ishikawa O, Ohigashi H, Imaoka S, Takahashi E, Nakamura Y. Related Articles, Nucleotide, OMIM, F
Isolation and characterization of a novel human pancreas-specific gene, pancpir, is down-regulated in pancreatic cancer cells.
Genes Chromosomes Cancer. 1998 Jul;22(3):179-85.
PMID: 9624529 [PubMed - indexed for MEDLINE]
- ☐ **13:** Caron PR. Related /
Computer analysis of cloned sequences.
Methods Mol Biol. 1997;69:247-60. No abstract available.
PMID: 9116857 [PubMed - indexed for MEDLINE]
- ☐ **14:** Hosokawa Y, Maeda Y, Seto M. Related Articles, Nucleotide, OMIM, F
Human Helios, an Ikaros-related zinc finger DNA binding protein: cDNA cloning and tissue expression pattern.
Immunogenetics. 1999 Oct;50(1-2):106-8. No abstract available.
PMID: 10541817 [PubMed - indexed for MEDLINE]
- ☐ **15:** Yamakawa T, Miyata S, Ogawa N, Koshikawa N, Yasumitsu H, Kanamori T, Miyazaki K. Related Articles, Nucleotide, F
cDNA cloning of a novel trypsin inhibitor with similarity to pathogenesis-related proteins, and its frequent expression in human brain cancer cells.
Biochim Biophys Acta. 1998 Jan 21;1395(2):202-8.
PMID: 9473672 [PubMed - indexed for MEDLINE]

- ☐ **16:** [Becker KG, Nagle JW, Canning RD, Dehejia AM, Polymeropoulos MH, Gado AM, Biddison WE, Drew PD.](#) Related Articles, Nucleotide, OMIM, F

Molecular cloning and mapping of a novel human KRAB domain-containing C2H2-type zinc finger to chromosome 7q36.1.

Genomics. 1997 May 1;41(3):502-4. No abstract available.

PMID: 9169157 [PubMed - indexed for MEDLINE]

- ☐ **17:** [Medlin JF.](#)

Related /

Fast-track cloning.

Environ Health Perspect. 1997 Dec;105(12):1310-1. No abstract available.

PMID: 9405327 [PubMed - indexed for MEDLINE]

- ☐ **18:** [Mason P, Bayol S, Loughna PT.](#)

Related /

The novel sarcomeric protein telethonin exhibits developmental and functional regulation.

Biochem Biophys Res Commun. 1999 Apr 21;257(3):699-703.

PMID: 10208846 [PubMed - indexed for MEDLINE]

- ☐ **19:** [Inoue S, Urano T, Ogawa S, Saito T, Orimo A, Hosoi T, Ouchi Y, Muramatsu M.](#) Related /

Molecular cloning of rat efp: expression and regulation in primary osteoblasts.

Biochem Biophys Res Commun. 1999 Aug 2;261(2):412-8.

PMID: 10425199 [PubMed - indexed for MEDLINE]

- ☐ **20:** [Kiess M, Scharm B, Aguzzi A, Hajnal A, Klemenz R, Schwarte-Waldhoff I, Schafer R.](#) Related Articles, Nucleotide, OMIM, F

Expression of ril, a novel LIM domain gene, is down-regulated in Hras-transfected cells and restored in phenotypic revertants.

Oncogene. 1995 Jan 5;10(1):61-8.

PMID: 7824279 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Sort	▼	Save	Text	Add to Clipboard	C
Show:	20	▼	Items 1-20 of 105	Page 1 of 6	Select page:	1	2	3

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Book
Search	PubMed	▼	for				Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		
Display		Abstract	▼	Sort	▼	Save	Text	Add to Clipboard

Entrez PubMed

☐ 1: J Cell Biol 2000 Oct 30;151(3):495-506 Related Articles, Nucleotide, Books, Li**FREE full text article at
www.jcb.org**

PubMed Services

**DRAL is a p53-responsive gene whose four and a half LIM dom:
protein product induces apoptosis.****Scholl FA, McLoughlin P, Ehler E, de Giovanni C, Schafer BW.**Division of Clinical Chemistry & Biochemistry, Department of Pediatrics, Univ
of Zurich, 8032 Zurich, Switzerland.

Related Resources

DRAL is a four and a half LIM domain protein identified because of its differer
expression between normal human myoblasts and the malignant counterparts,
rhabdomyosarcoma cells. In the current study, we demonstrate that transcription
the DRAL gene can be stimulated by p53, since transient expression of function
p53 in rhabdomyosarcoma cells as well as stimulation of endogenous p53 by io:
radiation in wild-type cells enhances DRAL mRNA levels. In support of these
observations, five potential p53 target sites could be identified in the promoter
of the human DRAL gene. To obtain insight into the possible functions of DRA
ectopic expression experiments were performed. Interestingly, DRAL expressio
efficiently triggered apoptosis in three cell lines of different origin to the extent
no cells could be generated that stably overexpressed this protein. However, tra
transfection experiments as well as immunofluorescence staining of the endoge:
protein allowed for the localization of DRAL in different cellular compartments:
namely cytoplasm, nucleus, focal contacts, as well as Z-discs and to a lesser ext
the M-bands in cardiac myofibrils. These data suggest that downregulation of D
might be involved in tumor development. Furthermore, DRAL expression might
important for heart function.

PMID: 11062252 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Add to Clipboard
---------	----------	---	------	---	------	------	------------------

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)

09428642

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:14:36 ON 17 SEP 2001

=> file medline biosis embase caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 09:14:46 ON 17 SEP 2001

FILE 'BIOSIS' ENTERED AT 09:14:46 ON 17 SEP 2001
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'EMBASE' ENTERED AT 09:14:46 ON 17 SEP 2001
COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 09:14:46 ON 17 SEP 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

=> s dral (s) androgen

L1 8 DRAL (S) ANDROGEN

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 2 DUP REM L1 (6 DUPLICATES REMOVED)

=> d l2 total ibib kwic

L2	ANSWER 1 OF 2	MEDLINE	DUPLICATE 1
ACCESSION NUMBER:	2001103482	MEDLINE	
DOCUMENT NUMBER:	20458893	PubMed ID: 11001931	
TITLE:	Alzheimer's disease-associated presenilin 2 interacts with DRAL, an LIM-domain protein.		
AUTHOR:	Tanahashi H; Tabira T		
CORPORATE SOURCE:	Division of Demyelinating Disease and Aging, National Institute of Neuroscience, 4-1-1 Ogawahigashi, Kodaira, Tokyo 187-8502, Japan.. tanahash@ncnp.go.jp		
SOURCE:	HUMAN MOLECULAR GENETICS, (2000 Sep 22) 9 (15) 2281-9. Journal code: BRC. ISSN: 0964-6906.		
PUB. COUNTRY:	ENGLAND: United Kingdom		
LANGUAGE:	English		
FILE SEGMENT:	Priority Journals		
ENTRY MONTH:	200102		
ENTRY DATE:	Entered STN: 20010322 Last Updated on STN: 20010322 Entered Medline: 20010208		
AB	Using the yeast two-hybrid system, we screened for proteins interacting with presenilin 2 (PS2) and cloned DRAL . DRAL is an		

LIM-only protein containing four LIM domains and an N-terminal half LIM domain. Previously **DRAL** has been cloned as a co-activator of the **androgen** receptor and as a protein interacting with a DNA replication regulatory protein, hCDC47. Our yeast two-hybrid assay showed that **DRAL** interacted with a hydrophilic loop region (amino acids 269-298) in the endoproteolytic N-terminal fragment of PS2, but not that of. . . this region, R275A, T280A, Q282A, R284A, N285A, P287T, I288L, F289A and S296A, in PS2 abolished the binding. This suggests that **DRAL** recognizes the PS2 structure specifically. The in vitro interaction was confirmed by affinity column assay and the physiological interactions between endogenous PS2 and **DRAL** by co-immunoprecipitation from human lung fibroblast MRC5 cells.

Furthermore,

in PS2-overexpressing HEK293 cells, we found an increase in the amount of **DRAL** in the membrane fraction and an increase in the amount of **DRAL** that was co-immunoprecipitated with PS2. The potential role of **DRAL** in the cellular signaling suggests that **DRAL** functions as an adaptor protein that links PS2 to an intracellular signaling.

L2 ANSWER 2 OF 2 MEDLINE DUPLICATE 2
 ACCESSION NUMBER: 2000120800 MEDLINE
 DOCUMENT NUMBER: 20120800 PubMed ID: 10654935
 TITLE: FHL2, a novel tissue-specific coactivator of the androgen receptor.
 AUTHOR: Muller J M; Isele U; Metzger E; Rempel A; Moser M; Pscherer
 CORPORATE SOURCE: A; Breyer T; Holubarsch C; Buettner R; Schule R
 Universitaets-Frauenklinik, Abteilung Frauenheilkunde und Geburtshilfe I, Klinikum der Universitaet Freiburg, Breisacherstrasse 117, 79106 Freiburg, Germany.
 SOURCE: EMBO JOURNAL, (2000 Feb 1) 19 (3) 359-69.
 Journal code: EMB; 8208664. ISSN: 0261-4189.
 PUB. COUNTRY: ENGLAND: United Kingdom
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200003
 ENTRY DATE: Entered STN: 20000327
 Last Updated on STN: 20000327
 Entered Medline: 20000310

AB . . . which nuclear receptor-cofactor interactions result in tissue-specific gene regulation are unclear. Here we characterize a novel tissue-specific coactivator for the **androgen** receptor (AR), which is identical to a previously reported protein FHL2/**DRAL** with unknown function. In the adult, FHL2 is expressed in the myocardium of the heart and in the epithelial cells. . .

=> d his

(FILE 'HOME' ENTERED AT 09:14:36 ON 17 SEP 2001)

FILE 'MEDLINE, BIOSIS, EMBASE, CAPLUS' ENTERED AT 09:14:46 ON 17 SEP 2001

L1 8 S DRAL (S) ANDROGEN
 L2 2 DUP REM L1 (6 DUPLICATES REMOVED)

=> s dral (s) transcrip?

L3 12 DRAL (S) TRANSCRIP?

=> dup rem l3

PROCESSING COMPLETED FOR L3

L4 6 DUP REM L3 (6 DUPLICATES REMOVED)

=> d l4 total ibib kwic

L4 ANSWER 1 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 1
ACCESSION NUMBER: 2001:71304 BIOSIS
DOCUMENT NUMBER: PREV200100071304
TITLE: Single nucleotide polymorphisms distinguish multiple
dopamine transporter alleles in primates: Implications for
association with attention deficit hyperactivity disorder
and other neuropsychiatric disorders.
AUTHOR(S): Miller, G. M.; De La Garza, R., II; Novak, M. A.; Madras,
B. K. (1)
CORPORATE SOURCE: (1) Division of Neurochemistry, Harvard Medical School,
NERPRC, One Pine Hill Drive, Southborough, MA, 01772-9102:
bertha_madras@hms.harvard.edu USA
SOURCE: Molecular Psychiatry, (January, 2001) Vol. 6, No. 1, pp.
50-58. print.
ISSN: 1359-4184.
DOCUMENT TYPE: Article
LANGUAGE: English
SUMMARY LANGUAGE: English

AB. . . tandem repeat (FNTR; 39 bases/12 repeats) was observed in all
animals. Accordingly, this FNTR is unbefitting an association of DAT
transcript length with hyperactivity. However, sequence analysis
revealed potential single nucleotide polymorphisms (SNPs), one of which
affects a Bst11071 restriction site.. . . hypothesis, we cloned a
portion of a novel 10-repeat allele from the human gene containing an SNP
that abolishes a **Dral** restriction site. We conclude that SNPs
create a diversity of DAT alleles between individuals that may be greater
than previously. . .

L4 ANSWER 2 OF 6 MEDLINE DUPLICATE 2
ACCESSION NUMBER: 2001042068 MEDLINE
DOCUMENT NUMBER: 20517437 PubMed ID: 11062252
TITLE: DRAL is a p53-responsive gene whose four and a half LIM
domain protein product induces apoptosis.
AUTHOR: Scholl F A; McLoughlin P; Ehler E; de Giovanni C; Schafer
B
CORPORATE SOURCE: W
Division of Clinical Chemistry & Biochemistry, Department
of Pediatrics, University of Zurich, 8032 Zurich,
Switzerland.
SOURCE: JOURNAL OF CELL BIOLOGY, (2000 Oct 30) 151 (3) 495-506.
Journal code: HMV. ISSN: 0021-9525.
PUB. COUNTRY: United States
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200012
ENTRY DATE: Entered STN: 20010322
Last Updated on STN: 20010322
Entered Medline: 20001207

AB **DRAL** is a four and a half LIM domain protein identified because
of its differential expression between normal human myoblasts and the
malignant counterparts, rhabdomyosarcoma cells. In the current study, we
demonstrate that **transcription** of the **DRAL** gene can be
stimulated by p53, since transient expression of functional p53 in
rhabdomyosarcoma cells as well as stimulation of endogenous p53 by
ionizing radiation in wild-type cells enhances **DRAL** mRNA levels.
In support of these observations, five potential p53 target sites could
be
identified in the promoter region of the human **DRAL** gene. To
obtain insight into the possible functions of **DRAL**, ectopic
expression experiments were performed. Interestingly, **DRAL**
expression efficiently triggered apoptosis in three cell lines of

different origin to the extent that no cells could be generated. . . this protein. However, transient transfection experiments as well as immunofluorescence staining of the endogenous protein allowed for the localization of **DRAL** in different cellular compartments, namely cytoplasm, nucleus, focal contacts, as well as Z-discs and to a lesser extent the M-bands in cardiac myofibrils. These data suggest that downregulation of **DRAL** might be involved in tumor development. Furthermore, **DRAL** expression might be important for heart function.

L4 ANSWER 3 OF 6 MEDLINE DUPLICATE 3
 ACCESSION NUMBER: 2000120800 MEDLINE
 DOCUMENT NUMBER: 20120800 PubMed ID: 10654935
 TITLE: FHL2, a novel tissue-specific coactivator of the androgen receptor.
 AUTHOR: Muller J M; Isele U; Metzger E; Rempel A; Moser M; Pscherer
 CORPORATE SOURCE: A; Breyer T; Holubarsch C; Buettner R; Schule R
 Universitats-Frauenklinik, Abteilung Frauenheilkunde und Geburtshilfe I, Klinikum der Universitat Freiburg, Breisacherstrasse 117, 79106 Freiburg, Germany.
 SOURCE: EMBO JOURNAL, (2000 Feb 1) 19 (3) 359-69.
 Journal code: EMB; 8208664. ISSN: 0261-4189.
 PUB. COUNTRY: ENGLAND: United Kingdom
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200003
 ENTRY DATE: Entered STN: 20000327
 Last Updated on STN: 20000327
 Entered Medline: 20000310

AB . . . Here we characterize a novel tissue-specific coactivator for the androgen receptor (AR), which is identical to a previously reported protein FHL2/**DRAL** with unknown function. In the adult, FHL2 is expressed in the myocardium of the heart and in the epithelial cells. . . binds specifically to the AR in vitro and in vivo. In an agonist- and AF-2-dependent manner FHL2 selectively increases the **transcriptional** activity of the AR, but not that of any other nuclear receptor. In addition, the **transcription** of the prostate-specific AR target gene probasin is coactivated by FHL2. Taken together, our data demonstrate that FHL2 is the. . .

L4 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2001 ACS
 ACCESSION NUMBER: 1999:10552 CAPLUS
 DOCUMENT NUMBER: 130:247523
 TITLE: Study of genetic polymorphism of Hungarian plum pox potyvirus isolates by RT-PCR method
 AUTHOR(S): Pribek, Dalma; Palkovics, L.; Gaborjanyi, R.
 CORPORATE SOURCE: Plant Protection Inst., Hung. Acad. Sci., Budapest, 1525, Hung.
 SOURCE: Novenyvedelem (Budapest) (1998), 34(11), 601-605
 CODEN: NVVDAW; ISSN: 0133-0829
 PUBLISHER: Agroinform Kiado es Nyomda
 DOCUMENT TYPE: Journal
 LANGUAGE: Hungarian

AB Fifteen representative samples were selected from more than one hundred plum pox potyvirus (PPV) isolates. We have previously demonstrated the existence of both M and D serotypes in Hungary by indirect ELISA (IDAS) using monoclonal antibodies. Some isolates represented intermediate serotypes. In this paper, a two step reverse **transcription** -polymerase chain reaction (RT-PCR) technique and digestion of the products with virus strain specific restriction enzymes (**Dral**, **Rsal**, **Sful**) was carried out to provide further evidence that both serotypes of PPV are common in Hungarian orchards.

L4 ANSWER 5 OF 6 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

ACCESSION NUMBER: 97300409 EMBASE
DOCUMENT NUMBER: 1997300409
TITLE: A major non-LTR retrotransposon of Bombyx mori, L1Bm.
AUTHOR: Ichimura S.; Mita K.; Sugaya K.
CORPORATE SOURCE: S. Ichimura, Division of Biology and Oncology, Natl. Inst.
of Radiological Sciences, Inage-ku, Chiba-shi 263, Japan
SOURCE: Journal of Molecular Evolution, (1997) 45/3 (253-264).
Refs: 23
ISSN: 0022-2844 CODEN: JMEVAU

COUNTRY: United States
DOCUMENT TYPE: Journal; Article
FILE SEGMENT: 022 Human Genetics
LANGUAGE: English
SUMMARY LANGUAGE: English

AB Repetitive sequences with oligo A tails were observed in **Dral** fragments of Bombyx mori genomic DNA. The full sequence of the element, an abundant non-LTR retrotransposon of B. mori, was determined by assembling inner restriction fragments. This element, designated L1Bm, contained two ORFs encoding a gag-like protein and reverse **transcriptase** (RT), respectively. An endonuclease domain was identified at the N-terminus of the RT sequence. The homology search of the amino. . .

L4 ANSWER 6 OF 6 MEDLINE

ACCESSION NUMBER: 96434502 MEDLINE
DOCUMENT NUMBER: 96434502 PubMed ID: 8837469
TITLE: Mapping of the ribosomal operons on the linear chromosomal DNA of Streptomyces ambofaciens DSM40697.
AUTHOR: Berger F; Fischer G; Kyriacou A; Decaris B; Leblond P
CORPORATE SOURCE: Laboratoire de Genetique et Microbiologie, Unite associee INRA 952, Faculte des Sciences, Universite Henri Poincare-Nancy 1, Vandoeuvre-les-Nancy, France.
SOURCE: FEMS MICROBIOLOGY LETTERS, (1996 Oct 1) 143 (2-3) 167-73.

Journal code: FML; 7705721. ISSN: 0378-1097.
PUB. COUNTRY: Netherlands
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199612
ENTRY DATE: Entered STN: 19970128
Last Updated on STN: 19970128
Entered Medline: 19961210

AB . . . genet internal transcribed spacer. The six rrn loci of S. ambofaciens were cloned as recombinant cosmids and located on the AseI-**Dral** physical map of the linear chromosomal DNA. For five of the six ribosomal gene sets, the **transcriptional** orientation was determined relative to the physical map and was shown to be divergent away from an oriC-like locus.

=> s dral (p) androgen

L5 8 DRAL (P) ANDROGEN

=> d his

(FILE 'HOME' ENTERED AT 09:14:36 ON 17 SEP 2001)

FILE 'MEDLINE, BIOSIS, EMBASE, CAPLUS' ENTERED AT 09:14:46 ON 17 SEP 2001

L1 8 S DRAL (S) ANDROGEN
L2 2 DUP REM L1 (6 DUPLICATES REMOVED)
L3 12 S DRAL (S) TRANSCRIP?
L4 6 DUP REM L3 (6 DUPLICATES REMOVED)

L5

8 S DRAL (P) ANDROGEN

=> log y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	21.43	21.64
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-0.59	-0.59

* * * * *

Dear valued customer,

Your feedback is important to us. Would you kindly take a moment to complete our survey? This survey will only take about 5-10 minutes to complete. Your responses will be kept confidential and will help us improve STN Express with Discover! for your future use. Please click on the following link to access the survey.

<http://www.cas.org/ONLINE/STN/ExpressSurveyForm.html?LOGINID=SSSPTA1649JXM>

* * * * *

STN INTERNATIONAL LOGOFF AT 09:19:17 ON 17 SEP 2001